

Unexpected behavior in ^3He films and the theory of fermions in two dimensions

Paulo F. Farinas

Instituto de Física Gleb Wataghin, Universidade Estadual de Campinas 13083-970, Campinas, São Paulo, Brazil
(February 1, 2008)

It is shown that the recently observed and unexplained behavior for the spin diffusion in ^3He films can be understood in terms of the Fermi liquid theory, provided nonperturbative effects characteristic of the two dimensions are taken into account. It is conjectured that such a behavior suggests the onset of a regime in which spin-current is conserved.

PACS numbers: 71.10.Ay, 67.80.Jd, 67.90.+z, 73.50.-h

The theory of Fermi liquids in two dimensions ($2d$) has been at the center of various settlements involving unexpected physics that has been arising in a broad range of interesting materials. Examples range from systems in the quantum Hall regime to copper-oxide superconductors, including quantum-fluid mixtures of ^3He - ^4He . These latter provided grounds for a recent observation of unexpected behavior, whose main cause will be argued in the following to be a consequence of the two dimensionality of a general system of fermions. In a recent Letter, [1] Sheldon and Hallock presented experimental results for spin diffusion in a thin film of ^3He on top of a superfluid ^4He film. The result for the spin-diffusion coefficient (D) as a function of the ^3He coverage (D_3), shows an upturn followed by a saturation and a downturn, as D_3 is increased. These features could not be reproduced by standard theoretical predictions. [2]

Throughout their past work, Hallock and collaborators have managed to develop a notable degree of control over the various regimes that may be set in the rich systems that constitute ^3He - ^4He films. [3] In the work reported in Ref. [1], care has been taken to guarantee that the ^3He film forms a $2d$ system of interacting fermions, with little influence from the ^4He superfluid film that seats below. Hence one should be able to account for the observed features using theoretical results from a $2d$ Landau theory.

It is argued in this article that the theory of $2d$ Fermi liquids *does* account for the observed behavior (regarding the existence of a peak in D as a function of D_3) provided one includes nonperturbative effects that are inherent to a Fermi liquid in two spatial dimensions. [4,5] It is also pointed that the onset of regimes in the $2d$ Fermi liquid for which spin-current is a conserved quantity, [6] is related to such a peaked behavior, however more experimental results are necessary to establish whether these regimes may be tuned or not.

I shall start with some remarks on existing theoretical results, and explain why the first attempts to fit the experimental data failed. I shall then devise the microscopic scenario that accounts for the physics underlying the peaked behavior observed, and calculate D , whose result is plotted in Fig.(1), compared to the experimental points. It should be stressed that the aim here is not to fully reproduce the experimental data, but yet uncover the physics that leads to the observed peak. Indeed, although the accuracy of the theoretical fitting is

quite remarkable and consistent with the long withstanding experimental observations of Owers-Bradley *et al.* [7] the existence of a saturation plateau is not accountable in the simple model studied here. One possible reason for such a behavior is discussed towards the end of the text.

The phenomenological result for the spin-diffusion coefficient in two dimensions was derived in Ref. [2] and reads

$$D = \frac{\hbar}{\pi m^*} \frac{(1 + F_0^a)^3 C(\lambda)}{(F_0^a)^2 (T_F/T)^2 \ln(T_F/T)}, \quad (1)$$

where $F_\ell^{a(s)}$ stands for the usual Landau antisymmetric (symmetric) parameters, m^* is the effective mass, T_F is the (Fermi) temperature, and $C(\lambda)$ is a function of all Landau parameters, given by Eq.(2). On fitting Eq.(1) to experimental data, one should be careful with the approximation that is taken for $C(\lambda)$. Based on the *strictly perturbative* results of Havens-Sacco and Widom, [8] one may reasonably disregard all high order Landau parameters, which leads to an approximately constant value for $C(\lambda)$. Hallock and Sheldon tried to fit their experimental results using this particular approach, with $C(\lambda) \sim 3/4$. However, the existence of nonperturbative effects intrinsic to the $2d$ Fermi liquid, predicted some time ago by Engelbrecht and Randeria, [4] suggests that such an approximation for the diffusion coefficient is not valid in general. Although single Landau parameters, which are present in the formulas of most the equilibrium properties, may not show any unusual behavior as a function of the density, the combination of various (usually infinite) Landau parameters may drive a $2d$ Fermi liquid into quite exotic regimes when the density is varied, as it has been pointed recently. [6,9]

To carry a more specific discussion, the full expression for $C(\lambda)$,

$$C(\lambda) = \frac{\lambda(A_0^a)^2}{4N^2(0)z^4} \sum_{n=1}^{\infty} \frac{(4n-1)/(2n^2-n)}{2n^2-n-1+\lambda|\Gamma_{\uparrow\downarrow\uparrow}^k(\pi)|^2}, \quad (2)$$

is conveniently written in terms of the vertex function of perturbation theory taken in the limit of zero frequency, [10] $\Gamma_{\sigma_1\sigma_2\sigma_3\sigma_4}^k(\theta)$, where θ is the angle between the momenta of two quasiparticles in a scattering process constrained by a Fermi circle and σ_i are spin labels. z and

$N(0)$ are respectively the Green's function residue and the density of states, both at the Fermi surface. Here λ stands for the inverse of a phase space sum, [11]

$$\lambda = \frac{8}{\sum_{\sigma_i; \theta=0, \pi} |\Gamma_{\sigma_1 \sigma_2 \sigma_3 \sigma_4}^k(\theta)|^2 \delta_{\sigma_1 + \sigma_2, \sigma_3 + \sigma_4}}.$$

It is a straightforward yet not pleasant task to obtain these expressions from the results of Ref. [2] This convenience of having closed expressions in terms of microscopic quantities is a remarkable amenability of the theory in two dimensions. The projection of $\Gamma_{\sigma_1 \sigma_2 \sigma_3 \sigma_4}^k(\theta)$ on a conveniently chosen basis defines the Landau scattering amplitudes $A_\ell^{a(s)}$. [12] Using the forward scattering sum rule for these amplitudes, $\sum_\ell A_\ell^a + A_0^s = 0$, one readily sees that if only the leading order $A_0^{a(s)}$ are kept then $C(\lambda)$ is constant. [13] Note, however, that varying the density may in principle drive the Landau coefficients to combine in such a way as to bring $\Gamma_{\uparrow\downarrow\uparrow}^k(\pi)$ down to small values, which causes D to increase. In this particular case, if the spin-diffusion relaxation time becomes very large, either another diffusion mechanism as spin-lattice relaxation sets in or, in the ideal $2d$ interacting liquid, the system will enter a regime for which spin-current is conserved and a spin-viscous damping mechanism becomes associated with the transport of magnetization throughout the sample. [6,9] A sharp raising of the spin-diffusion coefficient when one varies the density is then one of the signatures expected at the onset of such a regime.

In order to obtain a result that includes all the parameters, I start with the expression obtained by Engelbrecht, Randeria, and Zhang [14] for the vertex part in the limit of zero exchange momentum,

$$\Gamma_{\uparrow\downarrow\uparrow}^\omega(\theta) = \frac{\pi \hbar^2}{z^2 m_h} \left\{ 2\pi n a_s^2 \sin^2(\theta/2) + \frac{1}{\ln[2\pi n a_s^2 \cos^2(\theta/2)]} \right\}, \quad (3)$$

where m_h is the ^3He hydrodynamic mass, a_s is a hard-core potential radius, and n is the density, which is linearly related to the layer coverage D_3 in the experiment of Ref. [1]

The expansion parameter of the theory is $g = -(\ln 2\pi n a_s^2)^{-1}$, and for coverages ranging from 0 – 0.8, g falls within the interval 0 – 0.58. Equation (3) follows from an all-order ladder summation in the spirit of the long known results in three dimensions. [15] In this sense, both terms on the right side of Eq.(3) are nonperturbative. The first term arises from a pole in the vertex function below the particle-hole continuum, and is a particularity of two dimensions, since it results from the finite value of the density of states for small momenta. This pole is associated with a two-hole bound state. [4] Note that this pole term has an essential singularity in g and hence it cannot be determined to any finite order in perturbation theory. The second term contains the particle-hole

continuum contributions. Although these contributions are separable in the dilute limit defined by the small values of g , the existence of a pole also phase-shifts the continuum and on determining the unusual behavior in the spin-diffusion coefficient, *both* terms are necessary. However, the continuum alone yields no new physics in two dimensions, it is the counterpart of the known three dimensional rendered logarithmic divergences in the repulsive channel. [16] Indeed, a calculation using only the continuum (which will be shown elsewhere) leads to the usual behavior for D , consistent with the result of Ref. [8] in second order perturbation, and also with Eq.(1) for a constant $C(\lambda)$. This discussion establishes the two dimensional nonperturbative nature of the unusual behavior in D , as derived in what follows.




FIG. 1. Experimental data of Sheldon and Hallock (symbols) and the theoretical fits (solid lines) at the corresponding temperatures; same values of two fitting parameters are used for all temperatures: the effective hard-core radius $a_s = 0.75$ Å whilst the vertical amplitude is fixed using solely the first experimental point at 30mK.

In order to calculate the spin-diffusion coefficient from Eq.(1) knowledge of the zero frequency limit of the vertex part is necessary. These different limits of the vertex function are related to each other through a Bethe-Salpeter equation, [10]

$$\Gamma^k(\theta) = \Gamma^\omega(\theta) - N(0)z^2 \int_0^{2\pi} \frac{d\theta'}{2\pi} \Gamma^\omega(\theta - \theta') \Gamma^k(\theta'), \quad (4)$$

where the spin indices have been suppressed for convenience. Using Eq.(3) to define the kernel plus inhomogeneous term, this equation is numerically solved and the result used to calculate D from Eq.(1). The outcome is shown in Fig.(1) To fit the experimental data of Sheldon and Hallock at the given temperatures, a choice of

$a_s = 0.75 \text{ \AA}$ was made. Although the width of the peak is rather sensitive to this choice, it should be noted that this value for a_s is fairly close to the long withstanding theoretical fits to the magnetic susceptibility measurements of Owers-Bradley *et al.* in ^3He films on Grafoil. [7] Such fittings were achieved by using $a_s \sim 0.68 \text{ \AA}$ for a hard-core potential and $a_s \sim 0.73 \text{ \AA}$ for a 6-12 potential. [8] To account for the vertical scale, the theoretical result was multiplied by a constant fixed by making the first experimental point at $D_3 = 0.1$ for the 30 mK curve coincide with the theoretical value of D at the same coverage and temperature. The theoretical curves for all other temperatures follow *with no extra adjustments*. This fact, plus the consistency of the value for the hard-core radius with the experimental result for the magnetic susceptibility, which is an equilibrium response in the spin channel, are rather suggestive that the physics underlying the peaked behavior comes from the particularities of interacting fermions in a reduced geometry.




FIG. 2. Diffusion coefficient of Fig.(1) in a logarithmic scale.

The discontinuity in the derivative of D near $D_3 = 0.8$, and its nearly constant behavior for intermediate coverages are two features left unexplained by the above considerations. While the former is consistent with the promotion of ^3He atoms to the first excited surface state of the superfluid ^4He , as it is discussed by Sheldon and Hallock in Ref. [1] the latter needs further considerations: A full range plot of the theoretical result is shown in Fig.(2) for $T = 30 \text{ mK}$. The sharp peak shown in this figure is actually a divergence of D occurring for $D_3 \sim 0.3$ layers. This result holds for the *ideal* 2d Fermi liquid and indicates that a regime for which spin-current is conserved exists for the densities near the divergence. However, the quasiparticles are coupled to the Nuclepore substrate used in the experiment. This complication is not treated

here, and it causes the spin-diffusion to be several times smaller than the theoretical estimates (a mean field tortuosity factor, independent of the coverage, seems to be sufficient to correct the measured values). [1] Hence, the spin-diffusion cannot grow indefinitely in such a non-ideal system. The fact that for intermediate values of the coverage, D remains nearly constant as the coverage is varied seems to be consistent with the nearly constant “spin-lattice” relaxation times observed, suggesting that a saturation-like behavior might be coming from an underlying spin-relaxation mechanism with origin not in the 2d system of fermions but rather in the environment.

In summarizing, an explanation has been presented for the existence of a peak in the recent measurements of the spin-diffusion coefficient as a function of the density in ^3He films. Although the physical scenario builds from quite traditional results, it is conjectured that such a behavior might be signaling the existence of new regimes of the 2d Fermi liquids predicted recently for which spin-current is conserved. The experimental results are shown also to support the predictions of Engelbrecht and Randeria for the existence of a collective mode in systems of interacting fermions, which exists only in two dimensions.

The author acknowledges useful discussions with K.S. Bedell and C. Kübert, the hospitality of A. Caldeira at IFGW-UNICAMP where this work has been developed, and N. Studart for quickly bringing the online version of Ref. [1] to his attention. This work was supported in part by FAPESP, São Paulo, Brazil, Grant 98/07433-8.

-
- [1] P.A. Sheldon and R.B. Hallock, Phys. Rev. Lett. **85**, 1468 (2000).
 - [2] K. Miyake and W.J. Mullin, Phys. Rev. Lett. **50**, 197 (1983); J. Low Temp. Phys. **56**, 499 (1984).
 - [3] R.B. Hallock, Phys. Today, pg. 30 (June 1998) and references therein.
 - [4] J. Engelbrecht and M. Randeria, Phys. Rev. Lett. **65**, 1032 (1990).
 - [5] Two dimensions means here that the quasiparticle’s momenta are confined to a surface (the fermi gas seats on a surface state adsorbed on superfluid ^4He); this leads to a one dimensional Fermi surface.
 - [6] P.F. Farinas, K.S. Bedell, and N. Studart, Phys. Rev. Lett. **82**, 3851 (1999).
 - [7] J.R. Owers-Bradley, B.P. Cowan, M.G. Richards, and A.L. Thomson, Phys. Lett. **65A**, 424 (1978).
 - [8] S.M. Havens-Sacco and A. Widom, J. Low Temp. Phys. **40**, 357 (1980).
 - [9] P.F. Farinas, K.S. Bedell, and N. Studart, Physica B **280**, 93 (2000).
 - [10] A.A. Abrikosov, L.P. Gor’kov, and I.E. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Mechanics* (Dover, New York, 1975).
 - [11] Note that the quantity expressed here by λ differs from that of Ref. [2] (λ_-). They are related by $\lambda_- = 1 - \lambda |\Gamma_{\uparrow\downarrow\uparrow}^k(\pi)|^2$.

- [12] G. Baym and C. Pethick, *Landau Fermi-Liquid Theory* (Wiley, New York, 1991).
- [13] The precise value for this constant is easy to calculate: $C_0(\lambda) = \pi^2/12$, which is not much different from $3/4$.
- [14] J. Engelbrecht and M. Randeria, and L. Zhang, Phys. Rev. B **45**, 10135 (1992).
- [15] V.M. Galitskii, Zh. Eksp. Teor. Fiz. **34**, 151 (1958); Sov. Phys. JETP **7**, 104 (1958).
- [16] These are the known singularities that, in the attractive channel, signalize the breakdown of the Fermi liquid scenario towards a BCS state.